

# MATH 211

## Online Asynchronous Survey in Calculus and Analytical Geometry

Dr. Ahmed Kaffel

Department of Mathematical Sciences  
University of Wisconsin Milwaukee

Spring 2023



# Precise Definition of Limits

Recall the definition of limits:

Suppose  $f(x)$  is defined close to  $a$  (but not necessarily  $a$  itself). We write

$$\lim_{x \rightarrow a} f(x) = L$$

spoken: “the limit of  $f(x)$ , as  $x$  approaches  $a$ , is  $L$ ”

if we can make the values of  $f(x)$  arbitrarily close to  $L$  by taking  $x$  to be sufficiently close to  $a$  but not equal to  $a$ .

The intuitive definition of limits is for some purposes too vague:

- ▶ What means ‘make  $f(x)$  arbitrarily close to  $L$ ’ ?
- ▶ What means ‘taking  $x$  sufficiently close to  $a$ ’ ?

## Precise Definition of Limits: Example

$$f(x) = \begin{cases} 2x - 1 & \text{for } x \neq 3 \\ 6 & \text{for } x = 3 \end{cases}$$

Intuitively, when  $x$  is close to 3 but  $x \neq 3$  then  $f(x)$  is close to 5.

How close to 3 does  $x$  need to be for  $f(x)$  to differ from 5 less than 0.1?

- ▶ the distance of  $x$  to 3 is  $|x - 3|$
- ▶ the distance of  $f(x)$  to 5 is  $|f(x) - 5|$

To answer the question we need to find  $\delta > 0$  such that

$$|f(x) - 5| < 0.1 \quad \text{whenever} \quad 0 < |x - 3| < \delta$$

For  $x \neq 3$  we have

$$|f(x) - 5| = |(2x - 1) - 5| = |2x - 6| = 2|x - 3| < 0.1$$

Thus  $|f(x) - 5| < 0.1$  whenever  $0 < |x - 3| < 0.05$ ; i.e.  $\delta = 0.05$ .

## Precise Definition of Limits: Example

$$f(x) = \begin{cases} 2x - 1 & \text{for } x \neq 3 \\ 6 & \text{for } x = 3 \end{cases}$$

We have derived

$$|f(x) - 5| < 0.1 \quad \text{whenever} \quad 0 < |x - 3| < 0.05$$

In words this means:

If  $x$  is within a distance of 0.05 from 3 (and  $x \neq 3$ )  
then  $f(x)$  is within a distance of 0.1 from 5.

## Precise Definition of Limits: Example

$$f(x) = \begin{cases} 2x - 1 & \text{for } x \neq 3 \\ 6 & \text{for } x = 3 \end{cases}$$

Similarly, we find

$$|f(x) - 5| < 0.1 \quad \text{whenever} \quad 0 < |x - 3| < 0.05$$

$$|f(x) - 5| < 0.01 \quad \text{whenever} \quad 0 < |x - 3| < 0.005$$

$$|f(x) - 5| < 0.001 \quad \text{whenever} \quad 0 < |x - 3| < 0.0005$$

The distances  $0.1$ ,  $0.01$ ,  $\dots$  are called **error tolerance**.

## Precise Definition of Limits: Example

$$f(x) = \begin{cases} 2x - 1 & \text{for } x \neq 3 \\ 6 & \text{for } x = 3 \end{cases}$$

Similarly, we find

$$|f(x) - 5| < 0.1 \quad \text{whenever} \quad 0 < |x - 3| < \delta(0.1)$$

$$|f(x) - 5| < 0.01 \quad \text{whenever} \quad 0 < |x - 3| < 0.005$$

$$|f(x) - 5| < 0.001 \quad \text{whenever} \quad 0 < |x - 3| < 0.0005$$

The distances  $0.1$ ,  $0.01$ ,  $\dots$  are called **error tolerance**.

We have:  $\delta(0.1) = 0.05$

## Precise Definition of Limits: Example

$$f(x) = \begin{cases} 2x - 1 & \text{for } x \neq 3 \\ 6 & \text{for } x = 3 \end{cases}$$

Similarly, we find

$$|f(x) - 5| < 0.1 \quad \text{whenever} \quad 0 < |x - 3| < \delta(0.1)$$

$$|f(x) - 5| < 0.01 \quad \text{whenever} \quad 0 < |x - 3| < \delta(0.01)$$

$$|f(x) - 5| < 0.001 \quad \text{whenever} \quad 0 < |x - 3| < 0.0005$$

The distances  $0.1$ ,  $0.01$ ,  $\dots$  are called **error tolerance**.

We have:  $\delta(0.1) = 0.05$ ,  $\delta(0.01) = 0.005$

## Precise Definition of Limits: Example

$$f(x) = \begin{cases} 2x - 1 & \text{for } x \neq 3 \\ 6 & \text{for } x = 3 \end{cases}$$

Similarly, we find

$$|f(x) - 5| < 0.1 \quad \text{whenever} \quad 0 < |x - 3| < \delta(0.1)$$

$$|f(x) - 5| < 0.01 \quad \text{whenever} \quad 0 < |x - 3| < \delta(0.01)$$

$$|f(x) - 5| < 0.001 \quad \text{whenever} \quad 0 < |x - 3| < \delta(0.001)$$

The distances  $0.1$ ,  $0.01$ ,  $\dots$  are called **error tolerance**.

We have:  $\delta(0.1) = 0.05$ ,  $\delta(0.01) = 0.005$ ,  $\delta(0.001) = 0.0005$

Thus  $\delta(\epsilon)$  is a function of the error tolerance  $\epsilon$ !

We need to define  $\delta(\epsilon)$  for arbitrary error tolerance  $\epsilon > 0$ :

$$|f(x) - 5| < \epsilon \quad \text{whenever} \quad 0 < |x - 3| < \delta(\epsilon)$$

We want  $|f(x) - 5| = 2|x - 3| < \epsilon$ . We define  $\delta(\epsilon) = \epsilon/2$ .



## Precise Definition of Limits: Example

$$f(x) = \begin{cases} 2x - 1 & \text{for } x \neq 3 \\ 6 & \text{for } x = 3 \end{cases}$$

We define  $\delta(\epsilon) = \epsilon/2$ . Then the following holds

$$\text{if } 0 < |x - 3| < \delta(\epsilon) \quad \text{then} \quad |f(x) - 5| < \epsilon$$

In words this means:

If  $x$  is within a distance of  $\epsilon/2$  from 3 (and  $x \neq 3$ )  
then  $f(x)$  is within a distance of  $\epsilon$  from 5.

We can make  $\epsilon$  arbitrarily small (but greater 0),  
and thereby make  $f(x)$  arbitrarily close 5.

This motivates the precise definition of limits. . .

## Precise Definition of Limits

Let  $f$  be a function that is defined on some open interval that contains  $a$ , except possibly on  $a$  itself.

$$\lim_{x \rightarrow a} f(x) = L$$

if there exists a function  $\delta : (0, \infty) \rightarrow (0, \infty)$  s.t. for every  $\epsilon > 0$ :

$$\text{if } 0 < |a - x| < \delta(\epsilon) \quad \text{then} \quad |f(x) - L| < \epsilon$$

In words: No matter what  $\epsilon > 0$  we choose,  
if the distance of  $x$  to  $a$  is smaller than  $\delta(\epsilon)$  (and  $x \neq a$ )  
then the distance of  $f(x)$  to  $L$  is smaller than  $\epsilon$ .

We can make  $f$  **arbitrarily close** to  $L$  by taking  $\epsilon$  arbitrarily small.

Then  $x$  is **sufficiently close** to  $a$  if the distance is  $< \delta(\epsilon)$ .

## Precise Definition of Limits

Let  $f$  be a function that is defined on some open interval that contains  $a$ , except possibly on  $a$  itself.

$$\lim_{x \rightarrow a} f(x) = L$$

if there exists a function  $\delta : (0, \infty) \rightarrow (0, \infty)$  s.t. for every  $\epsilon > 0$ :

$$\text{if } 0 < |a - x| < \delta(\epsilon) \quad \text{then} \quad |f(x) - L| < \epsilon$$

The definition is **equivalent to the one in the book**:

$$\lim_{x \rightarrow a} f(x) = L$$

if for every  $\epsilon > 0$  there exists a number  $\delta > 0$  such that

$$\text{if } 0 < |a - x| < \delta \quad \text{then} \quad |f(x) - L| < \epsilon$$

# Precise Definition of Limits


$$\lim_{x \rightarrow a} f(x) = L$$

if for every  $\epsilon > 0$  there exists a number  $\delta > 0$  such that

$$\text{if } 0 < |a - x| < \delta \quad \text{then} \quad |f(x) - L| < \epsilon$$

## Geometric interpretation:

For any **small interval**  $(L - \epsilon, L + \epsilon)$  around  $L$ ,  
we can find **an interval**  $(a - \delta, a + \delta)$  around  $a$   
such that  $f$  maps all points in  $(a - \delta, a + \delta)$  into  $(L - \epsilon, L + \epsilon)$ .




# Precise Definition of Limits

$$\lim_{x \rightarrow a} f(x) = L$$

if for every  $\epsilon > 0$  there exists a number  $\delta > 0$  such that

$$\text{if } 0 < |a - x| < \delta \quad \text{then} \quad |f(x) - L| < \epsilon$$

**Alternative geometric interpretation:**



For every **interval  $I_L$**  around  $L$ ,  
find **interval  $I_a$**  around  $a$

such that

if we restrict the domain of  $f$  to  
 **$I_a$** , then the curve lies in  **$I_L$** .

# Precise Definition of Limits - Example

Proof that

$$\lim_{x \rightarrow 3} (4x - 5) = 7$$

Let  $\epsilon > 0$  be arbitrary (the error tolerance).

We need to find  $\delta$  such that

$$\text{if } 0 < |x - 3| < \delta \quad \text{then} \quad |(4x - 5) - 7| < \epsilon$$

We have

$$\begin{aligned} |(4x - 5) - 7| < \epsilon &\iff |4x - 12| < \epsilon \\ &\iff -\epsilon < 4x - 12 < \epsilon \\ &\iff -\frac{\epsilon}{4} < x - 3 < \frac{\epsilon}{4} \\ &\iff |x - 3| < \frac{\epsilon}{4} \end{aligned}$$

Thus  $\delta = \frac{\epsilon}{4}$ . If  $0 < |x - 3| < \frac{\epsilon}{4}$  then  $|(4x - 5) - 7| < \epsilon$ .

## Precise Definition of Limits - Example

If the next exam will be insanely hard,  
then many students will fail.

The words **if** and **then** are hugely important!

In exams many students write:

$$0 < |x - 3| < \frac{\epsilon}{4}$$

$$|(4x - 5) - 7| < \epsilon$$

which is wrong.

Correct is:

**If**  $0 < |x - 3| < \frac{\epsilon}{4}$

**then**  $|(4x - 5) - 7| < \epsilon$

## Precise Definition of Limits - Example

Find  $\delta > 0$  such that

$$\text{if } 0 < |x - 1| < \delta \quad \text{then} \quad |(x^2 - 5x + 6) - 2| < 0.2$$

Note that  $\delta$  is a bound on the distance of  $x$  from 1.

Lets say  $x = 1 + \delta$ . Then

$$\begin{aligned}(x^2 - 5x + 6) - 2 &= (1 + \delta)^2 - 5(1 + \delta) + 4 \\ &= (1 + 2\delta + \delta^2) - (5 + 5\delta) + 4 \\ &= \delta^2 - 3\delta\end{aligned}$$

Thus

$$|(x^2 - 5x + 6) - 2| < 0.2 \quad \iff \quad |\delta^2 - 3\delta| < 0.2$$

Assume that  $|\delta| < 1$  (we can make it as small as we want), then:

$$|\delta^2 - 3\delta| \leq |\delta^2| + |3\delta| \leq |\delta| + |3\delta| \leq 4|\delta|$$

Thus: if  $4|\delta| < 0.2$  then  $|(x^2 - 5x + 6) - 2| < 0.2$ .

Hence  $\delta = 0.04$  is a possible choice.



## Precise Definition of Limits: Example

Let  $\lim_{x \rightarrow a} f(x) = L_f$  and  $\lim_{x \rightarrow a} g(x) = L_g$ . Prove the sum law:

$$\lim_{x \rightarrow a} [f(x) + g(x)] = L_f + L_g$$

Let  $\epsilon > 0$  be arbitrary, we need to find  $\delta$  such that

$$\text{if } 0 < |x - a| < \delta \text{ then } |(f(x) + g(x)) - (L_f + L_g)| < \epsilon$$

Note that  $(f(x) + g(x)) - (L_f + L_g) = (f(x) - L_f) + (g(x) - L_g)$ .

We know that there exists  $\delta_f$  such that:

$$\text{if } 0 < |x - a| < \delta_f \text{ then } |f(x) - L_f| < \epsilon/2$$

and there exists  $\delta_g$  such that:

$$\text{if } 0 < |x - a| < \delta_g \text{ then } |g(x) - L_g| < \epsilon/2$$

We take  $\delta = \min(\delta_f, \delta_g)$ . If  $0 < |x - a| < \delta$  then

$$|f(x) - L_f| < \epsilon/2 \quad \text{and} \quad |g(x) - L_g| < \epsilon/2$$

and hence  $|(f(x) - L_f) + (g(x) - L_g)| < \epsilon$ .

# Precise Definition of One-Sided Limits

## Left-limit

$$\lim_{x \rightarrow a^-} f(x) = L$$

if for every  $\epsilon > 0$  there is a number  $\delta > 0$  such that

$$\text{if } a - \delta < x < a \text{ then } |f(x) - L| < \epsilon$$

## Right-limit

$$\lim_{x \rightarrow a^+} f(x) = L$$

if for every  $\epsilon > 0$  there is a number  $\delta > 0$  such that

$$\text{if } a < x < a + \delta \text{ then } |f(x) - L| < \epsilon$$

# Precise Definition of One-Sided Limits - Example

## Right-limit

$$\lim_{x \rightarrow a^+} f(x) = L$$

if for every  $\epsilon > 0$  there is a number  $\delta > 0$  such that

$$\text{if } a < x < a + \delta \quad \text{then} \quad |f(x) - L| < \epsilon$$

Proof that  $\lim_{x \rightarrow 0^+} \sqrt{x} = 0$ .

Let  $\epsilon > 0$ . We look for  $\delta > 0$  such that

$$\text{if } 0 < x < 0 + \delta \quad \text{then} \quad |\sqrt{x} - 0| < \epsilon$$

We have (since  $0 < x$ )

$$|\sqrt{x} - 0| = |\sqrt{x}| = \sqrt{x} < \epsilon \quad \implies \quad x < \epsilon^2$$

Thus  $\delta = \epsilon^2$ . If  $0 < x < 0 + \epsilon^2$  then  $|\sqrt{x} - 0| < \epsilon$ .

# Precise Definition of Infinite Limits

## Infinite Limit

$$\lim_{x \rightarrow a} f(x) = \infty$$

if for every positive number  $M$  there is  $\delta > 0$  such that

$$\text{if } 0 < |a - x| < \delta \quad \text{then } f(x) > M$$

## Negative Infinite Limit

$$\lim_{x \rightarrow a} f(x) = -\infty$$

if for every negative number  $M$  there is  $\delta > 0$  such that

$$\text{if } 0 < |a - x| < \delta \quad \text{then } f(x) < M$$

# Precise Definition of Infinite Limits - Example

## Infinite Limit

$$\lim_{x \rightarrow a} f(x) = \infty$$

if for every positive number  $M$  there is  $\delta > 0$  such that

$$\text{if } 0 < |a - x| < \delta \quad \text{then } f(x) > M$$

Proof that  $\lim_{x \rightarrow 0} \frac{1}{x^2} = \infty$ .

Let  $M$  be a positive number. We look for  $\delta$  such that

$$\text{if } 0 < |0 - x| < \delta \quad \text{then } \frac{1}{x^2} > M$$

We have:

$$\frac{1}{x^2} > M \iff 1 > M \cdot x^2 \iff \frac{1}{M} > x^2 \iff \sqrt{\frac{1}{M}} > |x|$$

Thus  $\delta = \sqrt{1/M}$ . If  $0 < |0 - x| < \sqrt{1/M}$  then  $\frac{1}{x^2} > M$ .