

1. This equation can be factored  $(x + 7)(x - 3) = 0$  so the roots are  $x_1 = -7$ ,  $x_2 = 3$ .
2. This equation can be factored  $(x - 8)(x - 1) = 0$  so the roots are  $x_1 = 1$ ,  $x_2 = 8$ .
3. This equation can be factored  $(2x - 9)(x + 2) = 0$  so the roots are  $x_1 = -2$ ,  $x_2 = \frac{9}{2}$ .
4. This equation cannot be factored, so we apply the quadratic formula and obtain complex roots:

$$x = \frac{-1 \pm \sqrt{1 - 20}}{2} = -\frac{1}{2} (1 \pm i\sqrt{19}).$$

5. This equation cannot be factored. so we apply the quadratic formula and obtain:

$$x = \frac{2 \pm \sqrt{4 + 28}}{2} = 1 \pm \sqrt{8}.$$

6. For the line, the  $y$ -intercept is  $(0, 2)$ , and the slope is  $m = 2$ . The  $x$ -intercept solves  $0 = 2x + 2$ , so  $x = -1$  and the  $x$ -intercept is  $(-1, 0)$ .

For the parabola, the  $y$ -intercept is  $(0, -5)$ . The  $x$ -intercepts solve


$$x^2 - 4x - 5 = (x - 5)(x + 1) = 0 \quad \text{or} \quad x = -1, 5.$$

Thus, the  $x$ -intercepts are  $(-1, 0)$  and  $(5, 0)$ . The  $x$  value of the vertex is the midpoint between the intercepts, so  $x_v = \frac{-1+5}{2} = 2$ . Since  $g(2) = -9$ , the vertex is  $(2, -9)$ .

The points of intersection satisfy  $f(x) = g(x)$  or  $2x + 2 = x^2 - 4x - 5$ , so

$$x^2 - 6x - 7 = (x + 1)(x - 7) = 0.$$


When  $x = -1$ , then  $f(-1) = 0$ , so this point of intersection is  $(-1, 0)$ . When  $x = 7$ , then  $f(7) = 16$ , so this point of intersection is  $(7, 16)$ . The graph is below



7. The height of the ball can be written in factored form

$$h(t) = -16t^2 + 48t + 64 = -16(t^2 - 3t - 4) = -16(t + 1)(t - 4)$$

From this factored form, it is easy to see that the ball hits the ground at  $t = 4$  sec. The vertex is the midpoint of the  $t$ -intercepts, so  $t = \frac{4-1}{2} = 1.5$  sec with  $h(1.5) = 100$  ft. Below is the graph of the height of the ball.



8. From the lecture notes, we have

$$[\text{H}^+] = \frac{1}{2} \left( -K_a + \sqrt{K_a^2 + 4K_a x} \right),$$

where  $x$  is the normality of the solution. For a 0.1N solution,  $x = 0.1$ , so

$$[\text{H}^+] = \frac{1}{2} \left( -1.75 \times 10^{-5} + \sqrt{(1.75 \times 10^{-5})^2 + 0.4(1.75 \times 10^{-5})} \right) = 0.001314.$$

It follows that the  $\text{pH} = -\log_{10} 0.001314 = 2.881$ .

Similarly, for a 1N solution,  $x = 1$ , so

$$[\text{H}^+] = \frac{1}{2} \left( -1.75 \times 10^{-5} + \sqrt{(1.75 \times 10^{-5})^2 + 4(1.75 \times 10^{-5})} \right) = 0.004175.$$

It follows that the  $\text{pH} = -\log_{10} 0.004175 = 2.379$ .



9. a. The perimeter of the rectangle is 40 cm, so  $2y + 2x = 40$ . It follows that  $y = 20 - x$ .

b. The area of a rectangle is  $A = xy$ , so

$$A(x) = 20x - x^2.$$

c. The domain of  $A(x)$  is  $0 < x < 20$ . The maximum area occurs at the vertex of the parabola shown below on the left, so  $x = 10$  and  $A_{max} = 100 \text{ cm}^2$ . Thus, the rectangle with the maximum area is a square.

d.  $A(x)$  is a parabola. The graph is shown below on the left.



10. a. The equilibrium population satisfies

$$g(P_e) = 0.02P_e - 0.000025P_e^2 = 0.02P_e(1 - 0.00125P_e) = 0,$$

so  $P_e = 0$  or  $P_e = \frac{1}{0.00125} = 800$  individuals.

b. The maximum growth rate occurs at the vertex of parabola, which satisfies  $P = 400$ , the midpoint between the  $P$ -intercepts (or equilibria) of the growth function. Thus, the maximum growth rate is  $g(400) = 0.02(400) - 0.000025(400)^2 = 4$  individuals per generation. A sketch of the graph  $g(P) = 0.02P - 0.000025P^2$  is shown above on the right.